

# SEASONAL CROP OUTLOOK

## Sorghum: February 2021

### SUMMARY


With the 2020/21 summer cropping season for north-eastern Australia (NEAUS) nearing completion the chances for an overall above average yielding sorghum crop have improved. However, areas planted to sorghum were lower than normal during the traditional planting window across southern QLD and NSW. There remains, however, large variation in the outlook among local regions. Most areas in QLD have sorghum yield expectations close to or slightly above the long-term median yield expectation. The exception is for some parts of southeast QLD (SEQ) and central QLD (CQ), which have yield expectations slightly below the long-term median yield. Most areas in northern and central New South Wales (CNSW, NNSW) have sorghum yield expectations above the long-term median yield for that region. Note: This yield outlook is based on an average of three plantings after a crop-free (fallow) period through the winter season and therefore areas with longer fallow practices are likely to have better yield prospects for the coming season.

### GENERAL CONDITIONS


Although highly variable, rainfall during January 2021 was average to above average rainfall across most parts of the NEAUS summer cropping region. Rainfall recorded during September to end of January was average across most of that region with the exception of parts of SEQ, which had below average rainfall for that period. Widespread above average rainfall is needed over the next month to induce late-planting, specifically in CQ where late plantings can occur until mid-February. The recent pattern of the SOI, i.e. “consistently positive”, at the end of January indicates an average chance of receiving above average rainfall for most of the summer grains cropping region over the next 3-months ([www.longpaddock.qld.gov.au](http://www.longpaddock.qld.gov.au)).

### OUTLOOK

This regional sorghum crop outlook is based on the assumption of cropping after winter fallow. The benchmark for this outlook is the simulated long-term median shire sorghum yield within the broad NEAUS cropping region (Map 1). The median yield is based on predicted performance over the past 120-years using an agro-climatic model for sorghum with long-term rainfall records (see descriptive note for more details). The percentage departure of the forecast median for this season from the long-term median shire sorghum yield is given in Map 2. Map 3 shows the current forecast shire median yield ranked relative to all years. Any areas coloured in light grey, yellow or red have a poor to very poor chance of having crops above the long-term median yield, whereas areas coloured in dark grey, green or blue have good to very good chances of producing higher yielding crops. Map 2 & 3 are derived by considering conditions up to date (end of January) and projecting forward based on rainfall conditions in years with SOI phase similar to this year - “consistently positive” in the December to January period. The calculation of benchmark yields and outlook chances do not take into account effects of poor crop nutrition or damage due to pests, diseases, frosts or extreme events (e.g. heat waves).



Map 1: Simulated long-term median shire yield derived from 1901 to 2019 climate using 2021 technology.



Map 2: Percentage departure of the current forecast median shire yield from the long-term shire median yield.

Yield outcomes vary geographically across most of the NEAUS summer cropping region. Map 2 shows that for this season, most areas in CQ, SWQ and NSW have positive forecast median yield deviations of 0 to 20% above the long-term median, while most areas in SEQ have negative forecast median yield deviations (-20% to 0) below the long-term median for that region. Furthermore, as shown in Map 3, almost all yield outcomes in NSW and SWQ are in the 60% to 80% percentile compared to all yield expectations over the last 120 years. In contrast, parts of CQ and most of SEQ are showing yield outcomes in the 20<sup>th</sup> to 40<sup>th</sup> percentile range. *Note: Final summer crop yield is usually more affected by in-crop rainfall and temperatures (during crop growth) than by the soil moisture at sowing, although this remains an important factor. The probability of yield outcomes presented here does not directly translate to total production figures.*

## POOR CROP CHANCE

At present, there are currently no shires with an increased chance for sorghum crop yield falling below the 10<sup>th</sup> percentile yield of all years (data not shown). It should be noted that these values are calculated as broad indicators for shire scale and do not apply to farm level.


## STATE OUTLOOK

The current regional outlook shows the forecast median yield for the entire NEAUS sorghum-cropping region on the 1<sup>st</sup> February is 3.28 t/ha, which is slightly above the long-term median of 2.95 t/ha (Graph A). There is however, a 10% chance that the state yield could be lower than 3.15 t/ha, or higher than 3.42 t/ha. At local regional level, Queensland (QLD), central Qld (CQ), southwest QLD (SWQ), southeast Qld (SEQ) and northern NSW (NNSW) (Map 1), the forecast yield (t/ha) ranges are as follows:


Region	Median (50%)	DFY (%)	Percentile (%)	Lt median
CQ	2.46	0	54	2.45
SEQ	3.71	3	56	3.62
SWQ	2.61	12	78	2.33
QLD	2.92	8	69	2.72
NNSW	3.95	11	88	3.55

\* DFY is the percentage departure of the forecast shire median yield from the long-term shire median (Lt median) sorghum yield.

With summer crops approaching maturity and harvest across most of the southern regions of NEAUS, CQ and SEQ regions have yield expectations close to the long-term regional sorghum yield expectation (i.e. 54<sup>th</sup> & 56<sup>th</sup> percentiles, respectively), while SWQ and NNSW have predicted medians in the top 25<sup>th</sup> percentile ranked relative to all years. The current SOI phase (“consistently positive”) indicates chances to receive above average rainfall are close to climatology (50:50) in most parts of NEAUS summer cropping region over the next 3-months. Widespread average rainfall is needed over the next couple of months to ensure good crop growth conditions especially around flowering for late-planted crops.



Map 3: Forecast median shire yield ranked relative to all years (%)



Graph A: NEAUS yield forecast trajectories (10<sup>th</sup>, 50<sup>th</sup> and 90<sup>th</sup> percentiles).

### DESCRIPTIVE NOTE:

The seasonal sorghum outlook is based on the integration of (i) a simple agro-climatic sorghum stress index model (i.e. Bare fallow routine - Ritchie, 1972; Sorghum stress index model adapted from - Fitzpatrick and Nix, 1969; Nix and Fitzpatrick, 1969), which is sensitive to water deficit or excess during the growing season, (ii) actual climate data up to the forecasting date and (iii) projected climate data after that date. These projected data are drawn from historical analogue years based on similarity to the prevailing phase of the Southern Oscillation Index (SOI) (Stone et al., 1996). The sorghum model was ran from 1 April the year before harvest in order to account for the influence of the winter fallow on starting soil moisture conditions. The model shire input parameters (i.e. plant available water content, planting rain & stress index period) have been selected based on the best fit when calibrated against actual shire sorghum yields from the Australian Bureau of Statistics (ABS) census years for the period 1983 to 2000, 2006, 2011, & 2016. Oz-Sorghum MII showed correlations (r) ranging from 0.62 to 0.93 within the main sorghum producing shires (35) of NE Australia. These shires contribute to 96% of total average production of all sorghum producing shires.