

# SEASONAL CROP OUTLOOK

## Sorghum: January 2022

### SUMMARY


The prospects for an above average sorghum crop for the entire Australian summer grain region for 2021/22 remain favourable. At this early stage, current soil water conditions and seasonal rainfall outlook indicate a very high chance for an above average yielding sorghum crop. There remains, however, some variation in the outlook among local regions. Most areas in central Queensland (CQ) have sorghum yield expectations slightly above the median, while most areas in southern QLD (SWQ & SEQ) and northern New South Wales (NNSW) have sorghum yield expectations above to very much above the long-term median yield. This crop outlook is based on a crop-free (fallow) period through the winter season and therefore areas with longer fallow practices are likely to have better yield prospects for the coming season. It should be noted, this is still early in the growing season, and the range of likely sorghum yield outcomes remains wide.

### GENERAL CONDITIONS


Rainfall during December was average to above average for most of the summer crop region. Current available soil moisture levels (simulated through winter fallow using APSIM) indicate near full soil moisture profiles across the entire NEAUS summer grains region. The recent pattern of the SOI, i.e., “consistently positive”, at the end of December indicates an increased chance of receiving above average rainfall for most of the summer grains cropping region over the next 3-months ([www.longpaddock.qld.gov.au](http://www.longpaddock.qld.gov.au)). Atmospheric indicators for ENSO are now showing a well-established “La Niña” system. This is likely to further increase the chances for a wetter than normal summer to autumn period. The impact will however vary across the region. Progress of the climate indicators such as the SOI and sea surface temperature anomalies can be followed here ([www.bom.gov.au/climate/enso](http://www.bom.gov.au/climate/enso)). Note: this outlook is only applicable to a short winter fallow cropping system (~9-month fallow period).

### OUTLOOK

The benchmark for this outlook is the simulated long-term median shire sorghum yield within the broad NEAUS cropping region (Map 1). The median yield is based on predicted performance over the past 121-years using an agro-climatic model for sorghum with long-term rainfall records (see descriptive note for more details). The percentage departure of the forecast median for this season from the long-term median shire sorghum yield is given in Map 2. Probability of exceeding the long-term shire median yield for this year is shown in Map 3. Any areas coloured in light grey, yellow or red have a poor to very poor chance of having crops above the long-term median yield, whereas areas coloured in dark grey, green or blue have good to very good chances of producing higher yielding crops. Map 2 & 3 are derived by considering conditions up to date (end of December) and projecting forward based on rainfall conditions in years with SOI phase similar to this year - “consistently positive” in the November to December period. The calculation of benchmark yields, and outlook chances do not take into account effects of poor crop nutrition or damage due to pests, diseases, frosts or extreme events (e.g. heat waves).



Map 1: Simulated long-term median shire yield derived from 1901 to 2020 using 2021 technology.



Map 2: Percentage departure of the current forecast median shire yield from the long-term shire median yield.

The current outlook combines effects of the recharge of soil moisture profiles and the current rainfall outlook, which is based on analogue years from history with the same SOI phase as at the end of December 2021. Map 2 shows that for this season, almost the entire region have positive forecast median yield deviations. Specifically, most areas of southern QLD and northern NSW have predicted yield deviations between 10 to 30% above the long-term median. Furthermore, as shown in Map 3, almost all yield outcomes have high chances of exceeding the long-term median yield for almost the entire NEAUS summer grain region (Map 3). At this early stage of the season, the range of likely yield outcomes for the 2021/2022 season (see Regional Outlook below) is still wide as much of the growing season remains in the projected forecast. Updating each month, as the season progresses, causes the range of yield outcomes to narrow towards the final realised yield at the end of the season. *Note: Final summer crop yield is usually more affected by in-crop rainfall and temperatures (during crop growth) than by the soil moisture at sowing, although this remains an important factor.*

## POOR CROP CHANCE

At present, this early in the growing season, no part of the entire summer cropping region has increased chance for sorghum crop yield to fall below the 10<sup>th</sup> percentile yield of all years (data not shown). It should be noted that these values are calculated as broad indicators for shire scale and do not apply to farm level.


## REGIONAL OUTLOOK

The current regional outlook shows a forecast yield of 3.32 t/ha with the entire forecast distribution now above the long-term simulated median yield (2.99 t/ha) for the NEAUS sorghum-cropping region (Graph A). There is however, a 10% chance that the state yield could be lower than 3.21 t/ha, or higher than 3.52 t/ha. At local regional level, Queensland (QLD), central Qld (CQ), southwest QLD (SWQ), southeast Qld (SEQ) and northern NSW (NNSW) (Map 3), the forecast yield (t/ha) ranges are as follows:


Region	Worst 10%	Median (50%)	Best (%)	Lt Median
CQ	2.42	2.65	2.79	2.47
SEQ	4.07	4.19	4.28	3.66
SWQ	2.53	2.77	2.99	2.38
QLD	2.99	3.15	3.31	2.75
NNSW	3.71	3.92	4.15	3.60

\*Lt Median: long-term median.

At this early stage of the season, all regions have yield expectation well above the long-term regional sorghum yield expectation. This is mainly a result of the extremely good start to the season in most parts and the well above average rainfall prospects during the next 3-months. However, a wide range of possible outcomes still exists. This range will narrow as the season progresses and the actual climate experienced is incorporated in the analysis. The current SOI phase of "consistently positive" indicates an increased chance to receive above average rainfall in most parts of the NEAUS summer cropping region over the next 3-months.



Map 3: Probability of exceeding the long-term simulated median shire sorghum yield.



Graph A: State level yield forecast trajectories (10<sup>th</sup>, 50<sup>th</sup> and 90<sup>th</sup> percentiles).

### DESCRIPTIVE NOTE:

The seasonal sorghum outlook is based on the integration of (i) a simple agro-climatic sorghum stress index model (i.e. Bare fallow routine - Ritchie, 1972; Sorghum stress index model adapted from - Fitzpatrick and Nix, 1969; Nix and Fitzpatrick, 1969), which is sensitive to water deficit or excess during the growing season, (ii) actual climate data up to the forecasting date and (iii) projected climate data after that date. These projected data are drawn from historical analogue years based on similarity to the prevailing phase of the Southern Oscillation Index (SOI) (Stone et al., 1996). The sorghum model simulated from 1<sup>st</sup> April the year before harvest to account for the influence of the winter fallow on starting soil moisture conditions. The model shire input parameters (i.e. plant available water content, planting rain & stress index period) have been selected based on the best fit when calibrated against actual shire sorghum yields from the Australian Bureau of Statistics (ABS) census years for the period 1983 to 2000, 2006, 2011, & 2016. Oz-Sorghum MII showed correlations (r) ranging from 0.62 to 0.93 within the main sorghum producing shires (35) of NE Australia. These shires contribute to 96% of total average production of all sorghum producing shires.